

# From Interstellar Ices to Polycyclic Aromatic Hydrocarbons

A symposium to honor Lou Allamandola's Contributions to the Molecular Universe

Annapolis, MD, USA - September 13<sup>th</sup> to September 17<sup>th</sup>, 2015

## Spatial studies of the PAH characteristics: how subtle variations reveal sub-populations with different molecular structure and charge.

Els Peeters<sup>1,2</sup>, David J. Stock<sup>1</sup>, Charles W. Bauschlicher<sup>3</sup>, Louis J. Allamandola<sup>4</sup>,  
Alexander G.G.M. Tielens<sup>5</sup>, Alessandra Ricca<sup>2</sup> and Mark G. Wolfire<sup>6</sup>

<sup>1</sup> Department of Physics and Astronomy, University of Western Ontario, London, ON N6A 3K7, Canada

<sup>2</sup> Carl Sagan Center, SETI Institute, 189 N. Bernardo Avenue, Suite 100, Mountain View, CA 94043, USA

<sup>3</sup> Entry Systems and Technology Division, Mail Stop 230-3, NASA Ames Research Center, Moffett Field, CA 94035

<sup>4</sup> NASA-Ames Research Center, Space Science Division, Mail Stop 245-6, Moffet Field, CA 94035, USA

<sup>5</sup> Leiden Observatory, PO Box 9513, 2300 RA Leiden, The Netherlands

<sup>6</sup> Astronomy Department, University of Maryland, College Park, MD 20742, USA

e-mail: epeeters@uwo.ca

The mid-IR spectra of photodissociation regions (PDRs) are dominated by the well-known emission features at 3.3, 6.2, 7.7, 11.3, and 12.7  $\mu\text{m}$ , generally attributed to polycyclic aromatic hydrocarbon molecules (PAHs). PAHs drive much of the physics and the chemistry in these PDRs, e.g. by heating the gas and as a catalyst in the formation of molecular hydrogen on their surfaces. Thus, PAHs and PDRs are intimately connected, and a complete knowledge of PDRs requires a good understanding of the properties of the PAH population.

One of the best ways to investigate the detailed characteristics of the PAH population is by analyzing IR spectral maps. Here, we present the results of such an analysis of spectral maps obtained with Spitzer/IRS in the 5-20  $\mu\text{m}$  range toward a sample of Galactic HII regions and reflection nebulae. These maps show subtle, but significant spatial variations in individual PAH emission bands. The overall dominant charge state of the PAH population is certainly a key factor in driving these variations. However, with our maps, we can also probe changes within the so-called ionic PAH emission bands that all originate from PAHs with the same charge state. We find that even within these bands, spatial variations occur that indicate contributions from at least 2 spatially distinct components, and thus PAH sub-populations, to the 7–9 micron PAH emission. We attribute these changes to molecular structure variations in the PAH sub-populations and discuss the astronomical implications.

## REFERENCES